

## Plasma levels of vitamin-D and C-reactive protein in adult Indian patients with COVID-19

Ravi Prakash<sup>1</sup>, R.B Singh<sup>2</sup>

<sup>1</sup> Department of Chemistry, B.S.A. College, Mathura, Uttar Pradesh, India

<sup>2</sup> Scientist, Department of Zoology, School of Life Sciences, Dr. Bhimrao Ambedkar University, Khandari Campus, Agra, Uttar Pradesh, India

### Abstract

Vitamin D deficiency (VDD) has been associated with increased mortality in patients with COVID-19. The anti-inflammatory properties of vitamin D are vital when considering the heightened inflammation in COVID-19 patients. Although avalanche of reports are available on the plasma levels of vitamin D and C-reactive protein (CRP) in COVID-19 patients, information on the dynamics of CRP level changes in different categories of vitamin D status is lacking. Therefore, the plasma levels of CRP were determined in COVID-19 patients with different vitamin D status. Forty-nine adults comprising 31 COVID-19 patients and 18 apparently healthy participants who served as controls were enrolled into this study. The plasma levels of high sensitivity-CRP (hsCRP) and vitamin D were determined using ELISA. Vitamin D status was classified as deficient, insufficient, sufficient and intoxication as appropriate. None of the COVID-19 patients had vitamin D insufficiency while majority of the controls were vitamin D sufficient. The proportion of COVID-19 patients with vitamin D intoxication was significantly higher than in the controls. The median plasma levels of vitamin D and hsCRP were not significantly different between the COVID-19 patients and the controls. The median plasma level of hsCRP was slightly lower in the vitamin D intoxication group compared with the vitamin D sufficient group. Vitamin D intoxication is common among Indian patients with COVID-19 and the vitamin D intoxication is associated with low plasma hsCRP level. There is the need for public enlightenment on the dangers inherent in vitamin D supplement abuse in Indian patients.

**Keywords:** Acute phase protein, Anti-inflammation, COVID-19, Vitamin D intoxication Indian patients

### Introduction

Coronavirus disease (COVID-19) is a global pandemic caused by the severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2). The high morbidity and mortality rate of the disease is associated with a number of factors [1]. These factors include presence of co-morbidities, nutritional deficiencies and gender among others [2, 3, 4, 5, 6]. Adequate and balanced nutrition are essential for proper body functioning and boosting of the immune system. Deficiencies of vitamins especially, vitamin D has been associated with increased mortality in critically ill patients [7].

Vitamin D is a pleiotropic hormone with critical role in maintenance of health [8]. It is involved in calcium and phosphorous homeostasis and also has antimicrobial and immunomodulatory properties [6]. In immune response, vitamin D plays important roles in the regulation of both the innate and adaptive immune system. Several cells of the immune system including neutrophil, monocytes, macrophages, B-cells and T-cells express vitamin D receptor (VDR) which directly or indirectly modulate their activities towards achieving immunity [9]. Vitamin D deficiency (VDD) is the most common vitamin deficiency which is seldom recognized even in the general population. Up to 50% of critically ill patients are vitamin D deficient and this has been shown to result in a 3-fold increase in mortality [10]. Vitamin D possesses anti-inflammatory and immune supporting properties and has thus gained tremendous attention during the COVID-19 pandemic. VDD has been shown to decrease the immune defences against COVID-19 and cause progression to severe disease. VDD was reported as the most prevalent micronutrient deficiency in COVID-19 patients as 76% and 24% of patients studied [11] had

moderate and severe VDD respectively. This observation is further elucidated by the reported inverse links between vitamin D and COVID-19 infection as well as outcomes [12, 13]. Similar reports have also shown that low vitamin D levels are associated with high mortality rate in COVID-19 patients [14, 15, 16].

The anti-inflammatory properties of vitamin D are vital when considering the heightened inflammation occasioned by cytokine storm in COVID-19 patients [17, 18]. One of the important acute phase proteins usually used as a biomarker for acute and chronic inflammation is C-reactive protein (CRP) [17]. It is a non-specific acute phase protein whose high levels and bimodal distribution have been used in triage, diagnostics and prognostication of COVID-19 patients [19, 20, 21], that CRP levels have a strong correlation with COVID-19 severity and prognosis. They showed that non-survivors had CRP levels about three-times higher than the levels in the survivors. Similarly, that plasma CRP level is a good predictor of COVID-19 severity and progression [22].

Although reports are abounding on the plasma levels of vitamin D and CRP in COVID-19 patients, information on the dynamics of CRP level changes in different categories of vitamin D status is lacking. This thus serves as the basis for this study. Recently the silico study on the inhibition of Corona virus disease (Covid-19) protease by Indian herbal plants [23].

### Materials and Methods

**Study Design:** This study was a case control study.

**Study Population:** A total of 49 participants were enrolled into this study. They comprised 31 patients with COVID-19

and 18 age and gender-matched apparently healthy participants who served as controls. All the COVID-19 patients were confirmed positive while the controls were certified negative using the real-time reverse-transcriptase polymerase-chain reaction (RT-PCR). The COVID-19 patients were followed up from the point of diagnosis till discharge from the Isolation Centre.

**Sample collection:** Venous blood (10 mL) was obtained from each SARS-CoV-2 infected patient upon diagnosis and at discharge and at enrolment only in controls. The blood samples were dispensed into lithium heparin containing sample bottles and plasma samples obtained were kept at -20 °C until analyzed.

**Laboratory Analyses:** The plasma levels of vitamin D and hsCRP were determined using ELISA following the Manufacturer's instruction JALMA Agra UP. (India)

**Vitamin D classification:** The plasma vitamin D level was Classified [24]. Vitamin D levels of < 10ng / mL, 10 - 30ng /mL, 30 - 100ng / mL and 100 ng/ml. were considered as deficiency, insufficiency, sufficiency and intoxication respectively.

**Statistical Analysis:** Statistical analysis was carried out using Mann Whitney U, Fisher's Exact test and Spearman

rank correlation as appropriate. P-values less than 0.05 were considered as statistically significant.

## Results and Discussion

The vitamin D status of the study participants are shown in Table 1. None of the study participants had vitamin D deficiency. Also, no COVID-19 Indian patient had vitamin D insufficiency whereas 11.1% of the controls had vitamin D insufficiency. The proportion of COVID-19 Indian patients with vitamin D intoxication was significantly higher than in the controls. Unlike the COVID-19 Indian patients, majority of the controls were vitamin D sufficient (Table-1).

**Table 1:** Distribution of the vitamin D status of the study participants. Data shown are frequency; n (%). in COVID-19 Indian Patients

Category	COVID-19	Control	N	P-value
Deficiency	0(0.0)	0(0.0)	0	0.038
Insufficiency	0(0.0)	2(11.1)	2	-
Sufficiency	15(48.4)	12(66.7)	27	-
Intoxication	16(51.6)	4(22.2)	20	-

\*Significant at P<0.05

The median levels of vitamin D were not significantly different between the COVID-19 Indian patients and the controls (P-value 0.059). Similarly, the median levels of hsCRP were similar between the 2 groups (P-value-0.173) and results are shown in Table-2.

**Table 2:** Baseline plasma levels of vitamin D and high sensitivity C-reactive protein (hsCRP) in COVID-19 Indian patients and the controls


Parameters	COVID-Diagnosis (n=31)	Controls (n=18)	P-value
Vitamin D(ng/ml)	100.39 (66.48 – 134.49)	72.49 (47.23 – 98.34)	0.059
hsCRP (mg/L)	1.56 (0.54 – 7.62)	2.09 (1.46 – 5.79)	0.173

There were no significant changes in the median plasma levels of vitamin D and hsCRP at diagnosis compared with the levels at discharge (Table-3).

**Table 3:** Plasma levels of vitamin D and hsCRP in COVID-19 Indian patients at diagnosis and at discharge

Parameters	COVID-Diagnosis (n=31)	Controls (n=31)	P-value
Vitamin D(ng/ml)	100.39 (66.48 – 134.49)	91.43 (68.56 – 124.02)	0.695
hsCRP (mg/L)	1.56 (0.54 – 7.62)	1.39 (0.63 – 5.77)	0.975

The correlation between vitamin D and hsCRP is shown in Figure-1. Vitamin D had an insignificant inverse correlation with CRP (r-value 0.095; P-value 0.471).



**Fig 1:** Correlation between the plasma level of vitamin D and hsCRP in

### COVID-19 Indian Patients

To further understand the relationship between the vitamin D status and the plasma levels of hsCRP in COVID-19 Indian patients, the hsCRP levels between the vitamin D sufficient and vitamin D intoxication groups were

**Table 4:** Plasma levels of CRP in different classes of vitamin D in COVID-19 Indian Patients

Parameter	Vitamin D sufficient (n=14)	Vitamin D intoxication (n=16)	P- value
hsCRP (mg/L)	1.65 (0.52 – 10.69)	1.29 (0.56 – 2.85)	0.633

Vitamin D and CRP play important but opposing roles in immune responses. Vitamin D possesses anti-inflammatory properties while the CRP is an acute phase reactant involved in the inflammatory responses. The observed high proportion of COVID-19 Indian patients with vitamin D intoxication is a source of concern. This observation might not be unconnected with indiscriminate use of various drugs and supplements such as vitamin D upon noticing signs and symptoms of SARS-CoV-2 infection by the study participants with COVID-19. This act could be occasioned by poor knowledge about COVID-19 and prevailing apprehensive situations with a view to boosting immunity against SARS-CoV 2 infection especially, during the early phase of the pandemic. Our observation is of public health importance as there is the need for public enlightenment on self-medication in future epidemics or pandemics. It is well known that vitamin D toxicity causes hypercalcemia, formation of calcium stones and kidney injury. The resulting hypercalcemia can also have profound effects on cardiac, renal and central nervous system functions [25,19]. Effects on the aforementioned organs can result in direct and/or indirect impacts on the immune response since physiological organ cross-talk is essential in maintaining equilibrium and homeostasis.

Reduced levels of vitamin D have been reported in COVID-19 Indian patients and deficiency has been associated with high mortality rate in the patients [12,13,14,15,16]. In this study, a contrast observation was made as there was no significant difference in the plasma levels of vitamin D in COVID-19 Indian patients compared with the controls. Our observation could be linked to the observed prevailing vitamin D intoxication in the studied COVID-19 Indian patients. This could also explain the observed similar levels of CRP between the two groups. Although reports have shown that COVID-19 is associated with hyperinflammation which is reflected in CRP elevation [19,21,22,26]. The COVID-19-associated hyperinflammation could have been masked by the expected, heightened anti-inflammatory activities of vitamin D occasioned by high proportion of vitamin D intoxication in COVID-19 patients.

Downward trend in CRP level has been reported in hospitalized patients with COVID-19 [25]. Similar observation was observed in this study as the median level of hsCRP was lower albeit insignificant; in COVID-19 patients at diagnosis compared with the level at discharge. This observation, in conjunction with other reports, further confirms the use of CRP in monitoring and prognostication of COVID-19 Indian Patients.

The observed inverse correlation between vitamin D level and hsCRP level is not surprising. This observation corroborates the anti-inflammatory and immune supporting properties of vitamin D which usually results in hyperinflammation dampening. These attributes of vitamin D could explain the observed lower median level of hsCRP

compared. It was observed that the median plasma level of hsCRP was nonsignificantly slightly lower in the vitamin D intoxication group compared with the vitamin D sufficient group (P-value 0.633) and results are shown as (Table-4).

in the vitamin D intoxication group compared with the vitamin D sufficient COVID-19 Indian patients.

It could be concluded from this study that vitamin D intoxication is common among Indian patients with vitamin D. Also, there was reduction in hsCRP level at discharge compared with the level at diagnosis in COVID-19 Indian patients. There is the need for public enlightenment on the dangers inherent in indiscriminate use of vitamin D as supplements by Indians as the expected benefit could be marred by the dangers associated with the toxicity of the drugs and supplements.

### References

- Richardson S, Hirsch JS, Narasimhan M, Crawford J M, McGinn T, Davidson KW, *et al.* Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City, Area', *Jama*,2020;323:2052-2059
- Rozenberg S, Vandromme J, Martin C. 'Are we equal in adversity. Does Covid-19 affect women and men differently, *Maturitas*,2020;138:62-68
- Velavan TP, Meyer CG. "Mild versus severe COVID-19: Laboratory markers, *Int J Infect Dis*, 2020,95,304-307
- Zhou F, Yu T, Du R, Fan G, Lin Y, Liu Z, *et al.* 'Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study', *Lancet*,2020;395:1054-1062(2018).
- Li G, Zhou C L, Ba Y M, Wang Y M, Song B, Cheng X B, *et al.* Nutritional risk and therapy for severe and critical COVID-19 patients: A multicenter retrospective observational study', *Clin Nutr*,2021,40,2154-2161
- Amrein K, Venkatesh B. 'Vitamin D and the critically ill patient', *Curr Opin Nutr Metab Care*,2012;15:188-193.
- Gombart AF, Pierre A, Maggini S, Alabyad D, Nahab FB, Creel-Bilos CM, *et al.* Jabaley CS. 'A Review of Micronutrients and the Immune System-Working in Harmony to Reduce the Risk of Infection', *Nutrients*,2020;12:236-240.
- Lee P. Vitamin D metabolism and deficiency in critical illness, *Best Pract Res Clin Endocrinol Metab*,2011;25:769-781
- Martens PJ, Gysemans C, Verstuyf A, Mathieu C. "Worsening severity of vitamin D deficiency is associated with increased length of stay, surgical intensive care unit cost, and mortality rate in surgical intensive care unit patients', *Am J Surg*,2020;204:37-43
- Flynn L, Zimmerman L H, McNorton K, Dolman M, Tyburski J, Baylor A, *et al.* Effects of vitamin D deficiency in critically ill surgical patients', *Am J Surg*, 2012,203,379-382
- Im JK, YS, Back J, Chung MH, Kwon HY, Lee JS, Nutritional status of patients with COVID-19', *Int J Infect. Dis*,2020;100:390-393.

12. Chandran M, Chan Maung A, Mithal A, Parameswaran R, 'Vitamin D in COVID-19: Dousing the fire or averting the storm? A perspective from the Asia-Pacific', *Osteoporos Sarcopenia*,2020;6:97-105.
13. Merzon E, Tworowski D, Garohovski, A., Vinker, S., Golan Cohen, A, Green I, Frenkel-Morgenstern M, Low plasma 25(OH) vitamin D level is associated with increased risk of COVID-19 infection: an Israeli population-based study,2020;287:3693-3702.
14. D'Avolio A, Avataneo V, Manca A, Cusato J, De Nicolò A, Lucchini R, *et al.* 25-Hydroxyvitamin D concentrations are lower in patients with positive PCR for SARS-CoV-2. *Nutrients*,2020;12(5):1359-1362.
15. Panagiotou G, Tee S A, Ihsan Y, Athar W, Marchitelli G, Kelly D, *et al.* Original publication: Low serum 25-hydroxyvitamin D (25(OH)D) levels in patients hospitalized with COVID-19 are associated with greater disease severity, *Clin Endocrinol (Oxf)*,2020;93:629-630
16. Shakoor H, Fechan J, Al Dhaheri A S, Ali H I, Platat C Ismail L C, Apostolopoulos V, *et al.* Immune-boosting role of vitamins D, C, E, zinc, selenium and omega-3 fatty acids: Could they help against COVID-19, *Maturitas*,2021;143:01-09
17. Arinola G O, Edem F V, Alonge T O. Levels of Plasma C-reactive protein, Albumin and Pre-Albumin in Indian COVID-19 Patients. *Annals of Medical Research*,2022;29(1):0046-0051.
18. Daneshkhah A, Agrawal V, Eshein A, Subramanian II, Roy. HK, Backman V. Evidence for possible association of vitamin D status with cytokine storm and unregulated inflammation in COVID-19 patients. *Aging Clin. Exp Res*,2020;32(10):2141-2158.
19. Lim K, Thadhani R 'Vitamin D Toxicity, *J Bras Nefrol*, 2020,42,238-44.
20. Lin F, Li L, Xu M, Wu J, Luo D, Zhu Y, *et al.* Prognostic value of interleukin-6, C-reactive protein, and procalcitonin in patients with COVID-19', *J Clin Virol*,2020;127:104370.
21. Stringer D, Braude P, Myint P K, Evans L, Collins J T, Verduri A, *et al.* The role of C-reactive protein as a prognostic marker in COVID-19, *Int J Epidemiol*,2020;50:420-429.
22. Ruan Q, Yang K, Wang W, Jiang L, Song J. 'Clinical predictors of mortality due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China', *Intensive Care Med*,2020;46:846-848.
23. Sadeghi-Haddad-Zavarch, Mahmoud Bayani, Masomeh Shokri, Mehran Ebrahimpour, Soheil Babazadeh, Arefch Mehracen, *et al.* Barary. Mohammad, and Hosseini, Akram. C-Reactive Protein as Prognostic Indicator in COVID-19 Patients, *Interdiscip Perspect Infect Dis*,2021;55:57582.
24. Prakash Ravi, Kumar Raj Singh RB, An in Silico study on the inhibition of Corona Virus Disease (COVID-19) Protease by the Indian herbal Plants, *International Journal of Pharmaceutical Research and Development*,2023;5(2):15-18.
25. Kader S, Comakli II, Tekindal MA, "Evaluation of serum vitamin D levels according to gender and age at Karapinar city: A follow-up study from Turkey', *Dubai Med J*,2019;2:141-145.
26. Sharifpour M, Rangaraju S, Liu M, Alabyad D, Nahab FB, Creel-Bulos C M, *et al.* protein as a prognostic indicator in hospitalized patients with COVID-19, *PLoS One*, 15: c0242400(2020).
27. Waterbury S. Implications of vitamin D toxicity & deficiency', *Nurse Pract*,2018;43:22-30.