



## Antimicrobial activity and characterization of bioactive compounds from flower extract of *Cassia javanica* Linn. Plant

R B Singh

Professor, Department of Zoology, School of Life Sciences, Dr. Bhimrao Ambedkar University, Khandari Campus, Agra, Uttar Pradesh, India

### Abstract

The present study was aimed to investigate the antimicrobial activity and characterization of bioactive constituents from flower extract of *Cassia javanica* Linn. Plant. Antimicrobial activity was tested against human pathogens in six different solvents using the well diffusion method. These, ethyl acetate and ethanolic crude extracts showed effective antimicrobial activity against all the tested human pathogens. The bioactive constituents of the flower extract revealed the presence of some biologically active substances like (alkaloids, cardiac glycosides, flavonoids, glycosides, phenolic compounds, phlobatannins, triterpenoids, saponins, volatile oils and tannins). In the GC-MS analysis, 14 bioactive compounds were identified. Present study concludes that the different extracts of *Cassia javanica* Linn. Flower contain a broad spectrum of bioactive compounds and also exhibit antimicrobial activity against all the tested human pathogens, therefore, *Cassia javanica* Linn. Flowers can be tapped source to discover the natural bioactive products that may serve as leads in the developments of novel therapeutic drugs.

**Keywords:** *Cassia javanica* Linn. flowers, antimicrobial activity, bioactive compounds, GC-MS

### Introduction

Plants have been a valuable source of natural products for maintaining human health, though the identification of active compounds from this valuable source came to light in recent years [1]. Medicinal plants constitute an important natural wealth of a country and they play a significant role in providing primary health care services to rural people [2] as well as substantial amount of foreign exchange can be earned by exporting medicinal plants to other countries [3]. Moreover, 80% of individual from developed countries use traditional medicines, having various compounds with medicinal properties [4].

A dramatic increase in microbial antibiotic resistance has developed over the last forty years in both the agriculture and medical sectors, this forced the researchers to develop new antimicrobial drugs for controlling pathogenic species [5]. In general, plants have the capacity to produce a large number of organic phytochemicals with complex structural diversity such as secondary metabolites [6,7], which served as plant self defense mechanisms against pests and pathogens [8] with interesting biological activities, this secondary metabolites can be utilized for medicinal purposes. Therefore, such plants should be investigated for better understanding their properties safety and efficiency [9]. Since plant products have no side effect [10] and they are comparatively cheaper. As a result, many researchers have paid attention to plants sources and biologically active compounds isolated from herbs [11],

The genus *Cassia* Linn. comprises 580 species of herbs, shrubs and trees, which are widely distributed throughout the world, of which only 20 species are indigenous to India [12]. Many of the *Cassia* spp. possess a good amount of medicinal properties [13] and they revealed potential in Ayurvedic medicine [14]. Especially, recent literature provides information about therapeutic uses of the plant *Cassia javanica* Linn. Bark of *Cassia javanica* Linn. is used as one of the ingredients in antidiabetic Ayurvedic

formulation. Leaves are proved to be active against Herpes simplex infection it is conventionally believed that this herb can reduce fever, decrease the virulence of pathogenic organisms, regulates the flow of chi and lubricate the intestine. In China it is applied to treat gastric pain, cold, malaria, measles, chickenpox and constipation [15].

*Cassia javanica* Linn. Plant is a small to medium-sized tree, upto 3-20 m tall with widely spreading horizontal branches and showy blossoms, trunk of young trees either smooth or armed with stump-remnants of branches. Leaves paripinnate with 5-15 pairs of leaflets. Taproot, petioles 1.5-4.0 cm long. Inflorescence an axillary racemose, many-flowered: flowers colour bright rose or pink, fading to white with red sepals (4-10 mm long), petals 15-35 mm long, whitish to reddish or buff, and stamens 10, 3 longer ones with filaments 2 cm long. 4 shorter with filaments about 1 cm long and 3 reduced with filaments about 1 cm long and minute anthers and phenology period is July to November. Since, very far of literature is currently available to substantiate antimicrobial prospective of the *Cassia javanica* Linn. Flower. The present attempt was made to examine the antimicrobial properties and bioactive compounds characterization of the *Cassia javanica* Linn. Flower.

### Materials and Methods

**Plant Material:** Fresh flowers of the species *Cassia javanica* Linn. (Family Caesalpiniaceae) were collected during January to April 2022 from, Forest Research Institute Dehradun (Uttarakhand). The flowers were separated from the plant and thoroughly washed with fresh water and kept for shade dry at room temperature to get rid off moisture, until further analysis.

**Preparation of extract:** Dried flower materials were powdered with electric blender, at room temperature and 2g of the powdered sample was soaked in 20 mL of different

solvents (ethanol, ethyl acetate, chloroform, benzene and water) overnight. Later, the samples were filtered under vacuum using Whatman No. 1 filter paper and stored in airtight screw-capped bottles at 5°C for further analysis.

**Preparation of Inoculum:** Seven clinical pathogenic organisms were obtained from the National JALMA Institute of Leprosy and other microbacterial Diseases Agra (India). Out of the seven organisms, four were *bacteria* *Escherichia coli*, *Salmonella typhi*, *Bacillus subtilis* and *Streptococcus pyogenes* and three were fungi *Candida albicans*, *Fusarium solani* and *Trichophyton rubrum*. Stock culture was maintained at 5°C on slopes of nutrient agar for bacteria and potato dextrose agar (PDA) for fungi. Active cultures for further experiments were prepared by transferring a loopful of cells from the stock cultures to test tubes of respective media. For antibacterial and antifungal activity, tubes were incubated for 24 h at 37 ± 2°C and 25 ± 2°C for 48 h. respectively. Muller-Hinton Broth (for

bacteria) and PD Broth (for fungi) were prepared for streaking and fresh slant cultures were prepared and stored in refrigerator at 5°C for future requirements.

**In vitro antimicrobial tests:** Spectrum of antibacterial activity was studied by using the techniques described [16]. Gentamicin sensitivity disc (30 mg: Hi-Media) was used as a positive control and respective solvents were taken as negative controls. At the end of incubation, inhibition zones formed around the discs were measured with transparent ruler in millimeter. These studies were performed in triplicate.

**Biochemical screening:** Biochemical tests in the flower of *Cassia javanica* Linn. Were done using standard methods: carotenoids [17], Total starch [18], total sugars [19], total proteins [20], total Frey amino acids [21], total phenols [22] and lipids [23]. Secondary metabolites were qualitatively tested by the standard methods [24, 25].

**Table 1:** Anitmicrobial activity of *Cassia javanica* Linn. Flower extracts in different silvents

S.No	Strains	Inhibition activity of <i>Cassia javanica</i> Linn. Flower extracts in different solvents.					
		A	B	C	ET	EA	Gentamicin
1	<i>Bacteria</i> <i>E. coli</i>	-	-	-	8.4±0.6	24.1±0.7	23.5
2	<i>S. typhi</i>	-	-	-	7.7±0.2	19.8±0.2	28.3
3	<i>B. subtilis</i>	11.5±0.4	11±0.25	-	10.2±0.4	21.8±0.4	25.8
4	<i>S. pyogenes</i>	-	-	-	8.6±0.05	13.7±0.8	31.7
5	<i>Fungi</i> <i>C. albicans</i>	18.6±0.02	11.6±0.07	15.8±0.6	18.2±0.2	21.3±0.2	35.6
6	<i>F. solani</i>	-	-	13.1±0.2	15.7±0.1	24.4±0.2	28
7	<i>T. rubrum</i>	-	-	-	-	19.7±0.4	26.2

Aqueous (A), Benzene(B), Chloroform(C), Ethanol (ET), Ethyl acetate (EA), No activity (-)

<sup>a</sup>- values are mean ± standard deviation of three determination

**GC-MS analysis:** GC-MS analysis of the *Cassia javanica* Linn. flower extract was performed using Perkin Elmer GC Clarus 500 MS system for identification of different components present in the extract, under the following conditions: column dimethyl polysiloxane DB-Ifused silica capillary column (30m x 0.25 mm x 0.1um of film thickness); carrier gas helium (1mL/min); injector temperature - 250 deg C; detector temperature - 200 deg C; column temperature - 35 - 180 deg \* C at 4 deg \* C / m \* in - 1 180 - 250 deg \* C 10° C/min; MS electron impact 70 V. Identification of the constituents was achieved based on comparison of mass spectra with the library ones (NIST Ver.2.1).

## Results and Discussion

Antimicrobial activities of *Cassia javanica* Linn. Flower extracts were screened against human pathogens (bacteria and fungi) using different solvents, viz. aqueous, benzene, chloroform, ethanol, and ethyl acetate. Among them, ethyl acetate and ethanol extract showed effective antimicrobial activity, In ethyl acetate, maximum inhibition was in *F. solani* (24.4 + 0.2mm) and the minimum in *S. pyogenes* (13.7 plus/minus 0.8 while in ethanol extract, the maximum inhibition was in *C. albicans* (18.2 plus/minus 0.2 mm) and the minimum in *S. typhi* (7.7 plus/minus 0.2 \* mm) Chloroform, benzene and aqueous showed moderate antimicrobial activity. The gentamicin (30 µg/dise). Was used positive control for comparing the bioassay (Table-1). Results of the biochemical screening in the fresh flower extract of *Cassia javanica* Linn. Were presented: carotenoids (0.731 plus/minus 0.03 \* mg / g) total soluble

sugars (37.83±6.64 mg/g), total soluble proteins (52.75 1.73 mg/g), total free amino acids (6.130±0.54 mg/g), total phenol (32.81±6.19 mg/g). Hydroxyl phenols (14.07± 4.69 mg/g) and lipids (13.33 5.77 mg/g). Further, alkaloids, cardiac glycosides, flavonoids, phenolic compounds, phlobatannins, tannins, tritrepnoides, volatile oils and saponins were also present in the *Cassia javanica* Linn. Flower.

Gas Chromatography and Mass Spectroscopy analyses were carried out on the flower extract of *Cassia javanica* Linn. And 14 compounds were identified. Active principles with their Retention time (RT), Molecular formula (MF), Molecular weight (MW), Concentration (%) and Nature of the compounds are presented in Table -1. The GC-MS chromatogram showed the peak identities of the identified compounds. In the present investigation. A variety of compounds has been detected viz. phenol, 4-propyl-(a). 3-O-methyl-d-glucose (b). Myo-Inositol, 4-C-methyl- (c), n-hexadecanoic acid (d). Hexadecanoic acid, ethyl ester (e), Phytol (1), 9-12-octadecadienoic acid (Z, Z) - (g), 9,12,15 octadecatrienoic acid, (Z,Z,Z) (h), 9.12-xtadecadienoic acid-methyl ester (E.E)- (1), octadecanoic acid (j). Octadecanoic acid-ethyl ester, phenol, 4-amino-3-methyl-(syn: m-cersol, 4-amino-), 1, 2-benzenedicarboxylic acid-diisooctyl ester (k) and propanal, 2-(4-ethoxyphenyl)-2-methyl.

Due to the value of medicinal plants as potential source of new compounds of therapeutic value researchers have been investigating them for their antimicrobial usefulness and as an alternative source to existing drugs [26,27], Plant extracts which are available for controlling microbial growth, non-

pollutive, cost effective, non hazardous and do not disturb ecological balance are preferred [28].

Different solvents have various degrees of solubility for different phytochemicals [29]. In present study, ethyl acetate and ethanol extracts had more inhibitory effects followed by chloroform, benzene and aqueous extracts and this study clearly revealed that organic solvents have more powerful antibacterial activity compared to water extracts [30, 31]. Previously, many researchers reported antimicrobial activity in *Cassia species*, viz. *Cassia nigricans* [32], *Cassia angustifolia* [33] and *Cassia fistula* Linn. [34, 35]. The present study also revealed a significant antimicrobial activity in *Cassia javanica* Linn. Flower extract.

Plant biochemicals are the vital source of innumerable number of antimicrobial compounds and several phytochemicals such as tannins, saponins, terpenoids, flavonoids, phenolics, polyphenols and alkaloids [36]. These compounds are potentially significant therapeutic application against human pathogens, including bacteria, fungi [37]. Scientist reported three flavonoids (5, 3 Dihydroxy-2 phenyl-4-chrome-7, 4-diglucoiside, Naringenin-7-glucoside and 3-glucoside of pelargonidin) from *Cassia javanica* Linn. Flower [38] reported alkaloids, flavonoids, carbohydrates, glycosides, triterpenoids, phenolic compounds [39] and saponins from *Cassia fistula* Linn. Which had interference with molecular targets in their pathogens. The major targets include biomembrane, proteins and nucleic acid and these are still regarded as a valuable pool for discovering novel mode of action [40].

To fulfill the needs of novel chemical substance, we require an in-depth knowledge of chemical constituents of plants [41] for application in pharmaceutical industry [42]. Therefore biological activities of 14 various compounds identified by GC-MS study of flower extract of *Cassia javanica* Linn. Were listed. Other species of genus *Cassia* reported as remedies to human diseases are: *Cassia auriculata* Linn. For the treatment of antihyperlipidaemic and antidiabetic [43]; *Cassia nigricans* can act as antifeedant, insecticide. Antiulcer, antiplasmodial, anti-inflammatory. Analgesic and antinociceptive [44]; *Cassia tora* Linn [45]. and *Cassia fistulal* Linn.. possess wound healing, ovicidal, antitumor, anti-inflammatory, antifungal, antibacterial and antioxidant activities and *Cassia italica* Lam. could be used as antioxidant, anti-inflammatory, antimicrobial, pesticide and cancer preventive [46]

## Conclusion

In conclusion, terrestrial plants have opened up a new perspective for pharmaceutical research and development of novel antimicrobial agents for the treatment of microbial diseases. In this study flower extracts of *Cassia javanica* Linn.. Demonstrated a broad spectrum of antimicrobial activity against human pathogens. Further, the present results supported its use as a traditional and folklore medicine. The bioactive compounds, present in *Cassia javanica* Linn. Flowers may lead to investigation of novel drugs against human diseases.

## References

1. Alam A, Sharma V, Sharma SC, Kumari P. Antibacterial activity of alcoholic extracts of *Entodon nepalensis* Mizush against some pathogenic bacteria. *Report Opin*,2012;4(10):44–47.
2. Rukia AK. *J Ethnobiol Ethnomed*,2007;3:1–4.
3. Phani Kumar G, Raj Kumar OP, Chaurasia, Singh SB. *J Med Plants Res*,2011;5(14):2929–2940.
4. Dahikar SB, Blutuda SA, Tambekar DH, Vibhute SK, Kashire SB. *Int J Pharm Sci Drug Res*,2011;3(1):32–34.
5. Daboor SM, Haroon AM. *Egypt J Aquat Res*,2012;38(4):233–239.
6. Kannan P, Ramadevi SR, Waheeta H. *Afr J Microbiol Res*,2009;3(4):180–184.
7. Gobalakrishnan R, Kulandarvelu M, Bhuvanewari R, Kandavel D, Kannan L. *J Pharm Anal*,2013;3(5):460–465.
8. Mazid M, Khan TA, Mohammad F. *Biol Med*,2011;3(2):232–239.
9. Nascimento GGF, Locatelli J, Freitas PC, Silva GL. *Braz J Microbiol*,2000;31(4):886–891.
10. Kavishankar GB, Lakshmidivi N, Mahadeva Murthy S, Prakash HS, Niranjana SR. *Int J Pharm Biomed Sci*,2011;2(3):65–80.
11. Essawi T, Srouf M. *J Ethnopharmacol*,2000;70(3):343–349.
12. CSIR. *The Wealth of India: A Dictionary of Indian Raw Materials*. New Delhi: Publications and Information Directorate,1950;11: 93–98.
13. Ganapaty S, Thomas PS, Ramana KV, Vidyadhar K, Chakradhar V. *J Nat Rem*,2002;2(2):102–120.
14. Ledwani L, Oberoi S. *Arch Appl Sci Res*,2010;2(1):295–301.
15. Chitam KP, Deore SL. *J Biomed Pharm Res*,2013;2(1):33–35.
16. Bauer AW, Kirby WM, Sherris JC, Turk M. *Am J Clin Pathol*,1966;45(4):493–495.
17. Goodwin TW, Britton G. *Distribution and analysis of carotenoids*. In: Goodwin TW, editor. *Plant Pigments*. London: Academic Press, 1988, 61–132.
18. McCready RM, Guggolz J, Silveira V, Owens HS. *Anal Chem*,1950;22(9):1156–1158.
19. Dubois M, Gilles KA, Hamilton JK, Rebers PA, Smith F. *Anal Chem*,1956;28(3):350–356.
20. Lowry OH, Rosenbrough NJ, Farr AL, Randall RJ. *J Biol Chem*,1951;193:265–275.
21. Troll W, Cannan RK. *J Biol Chem*,1953;200(3):803–811.
22. Swain T, Hillis WE. *J Sci Food Agric*,1959;10:63–68.
23. Jayaraman J. *Laboratory Manual in Biochemistry*. New Delhi: Wiley Eastern Ltd, 1981.
24. Harborne JB. *Phytochemical Methods: A Guide to Modern Techniques of Plant Analysis*. London: Chapman Hall, 1973, 49–188.
25. Odebiyi OO, Sofowora EA. *Lloydia*,1978;41(3):234–236.
26. Gowthami M, Tamilselvi S, Senthil Kumar, Panneerselvam A. *Asian J Plant Sci Res*,2012;2(2):110–114.
27. Kamble GS, Rasika CT, Pranav CC, Nirmala RD, Jyoti PS. *Pharma Biosci*,2011;2(2):197–201.
28. Joseph B, Muzafar AD, Vinod Kumar. *Glob J Biotechnol Biochem*,2008;2:56–59.
29. McCutcheon AR. *Clin Microbiol Rev*,1999;12(4):564–582.
30. Maji S, Dandapat P, Ojha D, *et al.* *J Phytol*,2010;2(4):57–64.
31. Vinoth B, Manivasagaperumal R, Balamurugan S. *Int J Res Biol Sci*,2012;2(3):98–101.
32. Ayo RG, Amupitan JO. *Chem Class J*, 2004, 24–26.

33. Makky EA, Masitah MY, Ibrahim MM. *Int J Chem Pharm Res*,2012;4(11):881–893.
34. Bhalerao SA, Kelkar TS. *Int Res J Biol Sci*,2012;1(5):79–84.
35. Panda SK, Padhi LP, Mohanty G. *J Adv Pharm Technol Res*,2011;2(1):62–67.
36. Alagesaboopathi C, Sivakumar R. *Int J Pharm Tech Res*,2011;3(1):27–31.
37. El-Astal ZY, Aera A, Aam A. *Pak Med Sci*,2005;21(2):187–193.
38. Pervez M, Obgeride ON, Irshad K, Ashfaq A. *Pak J Sci*,1995;47(1–2):34–37.
39. Pandya DJ, Patel VL, Desai TR, *et al.* *Int J Pharm Life Sci*,2012;3(2):1424–1429.
40. Yazdani D, Tan YH, Zainal Abidin MA, Jaganath I. *J Med Plants Res*,2011;5(30):6584–6589.
41. Yadav RNS, Agarwala M. *J Phytol*,2011;3(12):10–14.
42. Iqbal PJ. *Int J Drug Deliv Ther*,2012;2(4):135–138.
43. Yesu Raj J, Paul John Peter M, Joy V. *Asian J Plant Sci Res*,2012;2(2):187–192.
44. Ayo RG. *J Med Plants Res*,2010;4(14):1339–1348.
45. Choudhary M, Gulia Y, Nitesh. *Pharmacologyonline*,2011;3:78–96.
46. Sermakkani M, Thangapandian V. *Asian J Pharm Clin Res*,2012;5(2):90–94.